

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/21423>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

DNA VARIANTS

Leo A. J. Kluijtmans · Henk J. Blom · Godfried H. J. Boers
Bernard A. van Oost · Frans J. M. Trijbels
Lambert P. W. J. van den Heuvel

Two novel missense mutations in the cystathionine β-synthase gene in homocystinuric patients

Received: 12 December 1994 / Revised: 2 February 1995

Abstract Direct sequencing of the coding region of the cystathionine β-synthase (CBS) gene in two homocystinuric patients revealed the presence of two novel missense mutations. The first mutation, a 1111G → A transition, resulted in the substitution of the evolutionary conserved valine-371 by a methionine residue (V371M) and created a new *Nla*III restriction site. The second mutation, a G → A transition at base-pair 494, resulted in an amino acid change from cysteine to tyrosine (C165Y) and abolished a *Bso*FI restriction site. Both mutations were found in a compound heterozygous state with the previously described 833T → C transition.

Introduction

Homocystinuria caused by cystathionine β-synthase (CBS) deficiency is an inborn error of methionine metabolism and is inherited as an autosomal recessive trait. The predominant clinical manifestations of the homozygous state are arteriosclerosis, thromboembolic complications, skeletal abnormalities, ectopia lentis and mental retardation (Boers 1986; Mudd et al. 1989). The large range in the severity of the clinical features and in the biochemical response to therapy suggests genetic heterogeneity of the CBS gene. CBS, a pyridoxal-phosphate-requiring enzyme, catalyzes the synthesis of cystathionine from homocysteine and serine (Finkelstein and Martin 1984). To date,

14 mutations have been reported in the coding region of the CBS gene (Kraus 1994), which has been mapped to 21q22.3 (Müncke et al. 1988). In this report, we describe the identification of two new missense mutations in this coding region.

Materials and methods

Patient 1 was a 33-year-old man who suffered from ectopia lentis, osteoporosis, scoliosis and mental retardation. He also showed typically marfanoid features, such as arachnodactyly and dolichostenomelia. Therapy consisted of the daily administration of 750 mg pyridoxine, which reduced his total homocysteine levels to 35 μmol/l (control levels 8–18 μmol/l). Patient 2 was a 44-year-old male patient with many of the clinical features of homocystinuria: ectopia lentis, thrombosis, seizures, arachnodactyly and scoliosis. Family examination showed that a sister and a brother were also affected. All were pyridoxine-responsive.

For mutation detection, RNA was extracted from cultured fibroblasts (Chomczynski and Sacchi 1987) and first-strand cDNA was synthesized by standard methods. The cDNA product was amplified by the polymerase chain reaction (PCR) using three sets of oligonucleotides (Table 1). The generated PCR fragments covered the entire coding region and were subsequently subjected to direct sequencing according to the manufacturer's recommendations (Applied Biosystems, Forster City, CA, USA).

Table 1 Primers used for PCR amplification and sequence analysis of the cystathionine β-synthase cDNA

Primer	Nucleotide sequence (5' → 3')	Nucleotide position ^a
1 F	GTCAGCACCATCTGTCCGGTC	nt -26– -6
1 R	CTCGGAGCTCATCTTCTCTGGC	nt 528– 507
2 F	GATCGGGCTGGCCCTGGC	nt 453– 471
2 R	CCCGCCTCATCCACCACGG	nt 1358–1340
3 F	GACCAAGTTCCTGAGCGACAGGTG	nt 1146–1169
3 R	TCCTACCTGGCCGACTTCTCTC	nt 1720–1699

^aThe nucleotide position of the cystathionine-β-synthase cDNA is numbered according to Kraus et al. (1993)

L. A. J. Kluijtmans · H. J. Blom (✉) · J. M. F. Trijbels
L. P. W. J. van den Heuvel
Department of Paediatrics, University Hospital Nijmegen,
P.O. Box 9101, NL-6500 HB Nijmegen, The Netherlands

G. H. J. Boers
Department of Internal Medicine, Division of Endocrinology,
University Hospital Nijmegen, P.O. Box 9101,
NL-6500 HB Nijmegen, The Netherlands

B. A. van Oost
Department of Human Genetics, University Hospital Nijmegen,
P.O. Box 9101, NL-6500 HB Nijmegen, The Netherlands

Fig. 1 Comparison of amino acid sequences of homologous proteins of different species, *CysK-Ecoli Escherichia coli* acetylserine lyase; *M. circ.* CBS *Mucor circinelloides* cystathionine β -synthase. The amino acids are indicated by capital letters. A gap (–) has been introduced to maintain alignment

Human CBS:	E L Q E – G Q R C	V	V I L P D S V R
Rat CBS:	E L K E – G Q R C	V	V I L P D S V R
Yeast CBS:	E L T E – D D V I	V	A I F P D S I R
Wheat CS-precursor:	R P E N A G K L F	V	V V F P S F G E
<i>M. circ.</i> CBS:	E L K E – G Q R C	V	V I L P D S V R
<i>CysK-Ecoli:</i>		N K N I	V V I L P S S G E

Results and discussion

Direct sequence analysis of the entire coding region of the CBS gene in patient 1 showed a novel 1111G \rightarrow A transition and the previously described 833T \rightarrow C (I278T; Hu et al. 1993) missense mutation (data not shown). The observed G-to-A transition resulted in an amino acid exchange from valine to methionine at codon 371 of the mature protein (V371M) and introduced a new *Nla*III restriction site. Both transitions were confirmed at the genomic DNA level by *Nla*III and *Bsr*I restriction analysis, respectively. The substituted valine-371 residue appears to be highly conserved in homologous proteins in both prokaryotic and eukaryotic species (Fig. 1). Evolutionary conservation may indicate the functional relevance of an amino acid in the structure and function of the mature protein. Thus, this mutation is thought to interfere with the normal functioning of the CBS protein.

Patient 2 was found to be a compound heterozygote for the 833T \rightarrow C missense mutation and a novel 494G \rightarrow A transition. This latter transition resulted in a substitution of cysteine-165 by tyrosine (C165Y) in the CBS protein and abolished a *Bso*FI restriction site. The putative amine acid is located in the catalytic domain of the protein (J. P. Kraus, personal communications). This G-to-A transition may disrupt a disulphide bond in this catalytic domain. The disruption of a disulphide bond could interfere with intermolecular interactions and consequently with the catalytic activity of the protein. However, the present data provide no indications whether these two new missense mutations are compatible with pyridoxine responsiveness

in vivo, because the other mutation (833T \rightarrow C) is associated with responsiveness both in the homozygous and in the compound heterozygous states (Hu et al. 1993).

Acknowledgements The authors thank Hubert J.M. Smeets for his valuable discussion. This work was supported by research grant 93.176 from The Netherlands Heart Foundation.

References

- Boers GHJ (1986) Homocystinuria, a risk factor of premature vascular disease. *Clin Res Stud* 3:5–84
- Chomczynski P, Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156–159
- Finkelstein JD, Martin JJ (1984) Methionine metabolism in mammals. Distribution of homocysteine between competing pathways. *J Biol Chem* 259:9508–9513
- Hu FL, Gu Z, Kozich V, Kraus JP, Ramesh V, Shih V (1993) Molecular basis of cystathionine β -synthase deficiency in pyridoxine responsive and nonresponsive homocystinuria. *Hum Mol Genet* 2:1857–1860
- Kraus JP (1994) Molecular basis of phenotype expression in homocystinuria. *J Inher Metab Dis* 17:383–390
- Kraus JP, Le K, Swaroop M, Ohura T, Rosenberg LE, Roper MD, Kozich V (1993) Human cystathionine β -synthase cDNA: sequence, alternative splicing and expression in cultured cells. *Hum Mol Genet* 2:1633–1638
- Mudd SH, Levy HL, Skovby F (1989) Disorders of transsulfuration. In: Scriver CR, Beaudet AL, Sly WS, Valle D (eds) *The metabolic basis of inherited disease*, 6th edn. McGraw-Hill, New York, pp 693–734
- Müncke M, Kraus JP, Ohura T, Francke U (1988) The gene for cystathionine β -synthase (CBS) maps to the subtelomeric region on human chromosome 21q and to proximal mouse chromosome 17. *Am J Hum Genet* 42:550–559